


Base from U.S. Geological Survey Petersburg, 1960;  
Bradfield Canal, 1955; Sundum, 1961, 1971; Port  
Alexander, 1951, 1977; Sitka, 1951, 1970

SCALE 1:250000  
CONTOUR INTERVAL 200 FEET IN THE UNITED STATES AND 250 FEET IN CANADA  
DATUM IS MEAN SEA LEVEL



This report is preliminary and has not been  
reviewed for conformity with U.S. Geological  
Survey editorial standards.

**EXPLANATION**  
**SAMPLE SITES--Letters are explained on table 1.**

Anomalous site--sample locality at which the concentration is considered to deviate from the upper limit of normal background values, as determined by inspection of histograms, percentiles, and enrichment relative to crustal abundance.

**A Concentration**

**NOTE**

This map is one of a series of geochemical maps concerning the Petersburg area, southeast Alaska. For discussion of sample description, collection methods, media selection, sample preparation, statistical data, and analytical techniques, see Cathrall and others (1983).

**REFERENCE**

Cathrall, J. B., Day, G. W., Hoffman, J. D., and McDanal, S. K., 1983, A listing and statistical summary of analytical results for pebbles, stream sediments, and heavy-mineral concentrates from stream sediment, Petersburg area, southeast Alaska: U.S. Geological Survey Open-File Report 83-420-A.

**Table 1.--Nickel in 1430 nonmagnetic samples of heavy-mineral concentrates from stream sediments, Petersburg area, southeast Alaska.**

[Concentrations in parts per million; <, detected, but less than value shown; N, not detected at limit of detection or at value shown.  
Arithmetic mean, 27.7; standard deviation, 53; geometric mean, 15.8; and geometric deviation, 2.3, based on unqualified values within the sample population.]

Concentration	Map symbol	Frequency	Percentile
1,000	A	1	100
700	B	0	99.93
500	C	5	99.93
300	D	2	99.58
200	F	13	99.44
150	H	31	98.53
100	J	39	96.36
70	K	82	93.64
50	L	73	87.90
30	M	38	82.80
20	N	64	80.14
15	O	2	75.66
10	P	980	75.52
<10	R	14	6.99
N10	X	86	6.01
			0.00

**DISTRIBUTION AND ABUNDANCE OF NICKEL, DETERMINED BY SPECTROGRAPHIC ANALYSIS, IN NONMAGNETIC FRACTION OF HEAVY-MINERAL CONCENTRATES FROM STREAM SEDIMENTS, PETERSBURG AREA, SOUTHEAST ALASKA**

**By**

John B. Cathrall, Gordon W. Day, James D. Hoffman,  
and Steven K. McDanal  
1983